

Document downloaded from:

<http://hdl.handle.net/10251/150315>

This paper must be cited as:

Jiménez Molero, MC.; Miranda Alonso, MÁ.; Vayá Pérez, I. (2005). Triplet excited states as chiral reporter for the binding of drugs to transport proteins. *Journal of the American Chemical Society*. 127(29):10134-10135. <https://doi.org/10.1021/ja0514489>



The final publication is available at

<https://doi.org/10.1021/ja0514489>

Copyright American Chemical Society

Additional Information

Triplet excited states as chiral reporters for the binding of drugs to transport proteins

M. Consuelo Jiménez, Miguel A. Miranda* and Ignacio Vayá

Departamento de Química-Instituto de Tecnología Química UPV-CSIC, Universidad Politécnica de Valencia, camino de Vera s/n, E-46022 Valencia, Spain.

RECEIVED DATE (automatically inserted by publisher); e-mail: mmiranda@qim.upv.es

Human serum albumin (HSA) is the most important transport protein. One of its main functions is to carry different endogenous and exogenous agents (including drugs) around the blood-stream, in order to achieve their efficient and selective delivery to the specific targets.¹

The pioneering work performed by Sudlow and coworkers three decades ago revealed that there are two major and structurally selective binding sites in HSA, namely site I and site II.² Although more recent studies have found evidence supporting the existence of several subdomains, consensus still exists that there are basically two high affinity binding sites for small heterocyclic or aromatic compounds in this protein.³

As the knowledge of drug-HSA interactions is essential for understanding the biodistribution, metabolism, elimination and pharmacological effect of drugs in the body, a number of techniques have been used to address this problem. They include equilibrium dialysis, ultrafiltration, ultracentrifugation, calorimetry, fluorescence quenching, circular dichroism, liquid chromatography, capillary electrophoresis, NMR diffusion measurements, etc.^{4,5} Each method presents advantages and limitations; the latter are mainly related to sensitivity, interferences, diffusion problems or lack of reproducibility arising from a complicated workup.

Arylpropionic acid derivatives are the main group of non-steroidal anti-inflammatory drugs (NSAIDs). They are currently marketed as racemic mixtures (with few exceptions such as naproxen); however, their anti-inflammatory activity is mainly attributed to the (*S*)-isomer.⁶ Those drugs bind preferentially (and often stereoselectively) to site II of HSA by means of hydrogen bonding and electrostatic interactions.⁷ Upon methylation of the carboxy group, binding occurs preferentially to site I, where hydrophobic interactions predominate.⁸

The aim of the present work has been to explore the suitability of triplet excited states as chiral reporters for the important issue of drug binding to transport proteins. As the properties of these states are very sensitive to the experienced microenvironment,⁹ they should allow to study the drug distribution among the bulk solution and the different protein binding sites. More interestingly, triplet lifetime measurements would provide valuable information on the nature of drug-HSA complexes, such as strength of the interaction, conformational restrictions, protection from attack by oxygen or other reagents, stereochemical requirements, etc. Previously, laser flash photolysis (LFP) has been employed for the study of carprofen-HSA complexes.¹⁰ However, in this case fast reaction of the drug triplet state with the Trp unit of site I limits its possibilities to be used as a probe. In order to circumvent this problem, an “inert” triplet would be more adequate as reporter, while residing as observer in the HSA binding sites. Flurbiprofen (FBP) is a NSAID of the 2-arylpropionic acid class used for several medical purposes, such as the treatment of osteoarthritis, rheumatoid

arthritis, post-operative dental pain and ocular disorders.¹¹ The lowest triplet excited state of FBP has recently been characterized,¹² and its lifetime has been found to be highly sensitive to the medium. In the present work, the triplet state of FBP methyl ester (FBPMe, Chart 1) has been used as probe for the binding sites of HSA. The choice of FBPMe has been done for experimental convenience, since its hydrophobic character should favor inclusion within HSA (preferentially in site I, as stated above). Both (*R*)- and (*S*)-FBPMe have been investigated in order to obtain valuable information about the possible stereodifferentiation in the interaction of drugs with proteins.




Chart 1

Aqueous solutions containing (*S*)-FBPMe and HSA (molar ratio from 10:1 to 0.5:1) were prepared in neutral buffer (0.01 M PBS) and submitted to laser flash photolysis. The transient absorption spectrum obtained after laser excitation ($\lambda_{\text{exc}} = 266$ nm) of an aerated $2.5 \cdot 10^{-5}$ M solution of (*S*)-FBPMe displayed a maximum at 360 nm, assigned to the first triplet excited state.¹² Its decay (Figure 1) followed a first order exponential law with a lifetime of 1.5 μs .




Figure 1. Laser flash photolysis ($\lambda_{\text{exc}} = 266$ nm) of (*S*)-FBPMe (\blacktriangle), (*S*)-FBPMe/HSA 2:1 (\blacktriangle) and (*S*)-FBPMe/HSA 0.5:1 (\blacktriangle) in 0.01 M PBS. Normalized decays monitored at 360 nm. The concentration of (*S*)-FBPMe was $2.5 \cdot 10^{-5}$ M in all cases.

Remarkably, in the presence of HSA this species lived much longer (see Figure 1) and did not exhibit a first order exponential decay. Thus, when the [*S*]-FBPMe]/[HSA] ratio was between 1:1 and 0.5:1, the signal followed a second order exponential decay with lifetimes of 4.1 and 31.5 μs . The negligible contribution of the 1.5 μs lifetime component indicated the absence of free (*S*)-FBPMe in solution under these conditions.

It is noteworthy that protein-bound (*S*)-FBPMe has dramatically longer lifetimes than the non-complexed form (τ_{T} values up to 21 fold higher). This can be attributed to a slower

deactivation *via* non-radiative processes, due to important restrictions in the degrees of freedom inside the HSA binding sites, where a microenvironment is provided that protects the triplet excited state from attack by oxygen or other reagents. In this context, the fact that two different τ values were measured in the presence of HSA can be correlated with the existence of two types of binding sites. At higher $[(S)\text{-FBPMe}]/[\text{HSA}]$ ratios (from 1.25:1 to 10:1), three lifetimes values (1.5, 4.1 and 31.5 μs) were necessary to obtain a good fitting of the decay signal (also shown in Figure 1). This can be clearly associated with the presence of free (S)-FBPMe in solution, in addition to the two HSA-bound species.

Parallel experiments were carried out for (R)-FBPMe. In PBS solution the transient absorption spectra and the triplet lifetimes did not differ from those of the (S)- enantiomer. By contrast, when included within HSA, two components with τ 157.6 μs and 16.6 μs were found for (R)-FBPMe. These values are much higher than those obtained for (S)-FBPMe, indicating a remarkable stereodifferentiation between the two enantiomers in the protein binding process. A comparison between the decays monitored at 360 nm for (S)-FBPMe, (S)-FBPMe/HSA and (R)-FBPMe/HSA is shown in Figure 2.




Figure 2. Laser flash photolysis ($\lambda_{\text{exc}} = 266 \text{ nm}$) of (S)- or (R)-FBPMe $2.5 \cdot 10^{-5} \text{ M}$ (●), as well as 1:1 solutions of (S)-FBPMe/HSA (●) and (R)-FBPMe/HSA (●) in 0.01 M PBS, under air. **A:** Spectra obtained 3.7 μs after the laser pulse. **B:** Decays monitored at 360 nm.

Regression analysis of the decay curves for several $[\text{FBPMe}]/[\text{HSA}]$ ratios was performed to obtain the values of the A_1 , A_2 and A_3 coefficients (components with different lifetimes). Based on the known fact that the high affinity site of 2-arylpropionic acids methyl esters is site I,⁸ the major (longer-lived) component under non-saturating conditions was assigned to FBPMe within this site; while the minor (shorter-lived) component was assigned to site II-bound FBPMe. In order to obtain independent evidence supporting this assignment, additional LFP experiments were performed in the presence of warfarin, a well established probe for site I.¹³ In agreement with our initial assumption, addition of this probe resulted in a reduced contribution of the longer lifetime components (attributed to site I). Thus, at 1:1.1 $[\text{FBPMe}]/[\text{HSA}]/[\text{warfarin}]$ molar ratio this value dropped to *ca.* 50% (as compared with *ca.* 70% in the absence of warfarin).

The trends clearly show (Figure 3A) that, for the same $[\text{FBPMe}]/[\text{HSA}]$ ratio, inclusion within HSA is slightly favored for the (S)- isomer. Besides, a significant stereodifferentiation is observed when comparing the occupation level of the two sites, with higher preference for site I in the case of (S)-FBPMe. In general, when the $[\text{FBPMe}]/[\text{HSA}]$ ratio is low (far from saturation) all the drug is protein-bound and the high affinity site I is selectively populated. Progressive increase of the $[\text{FBPMe}]/[\text{HSA}]$ ratio results in saturation of site I and

(subsequently) site II. This leads to a decrease of site selectivity (Figure 3B) and the appearance of increasing amounts of free FBPMe.





Figure 3. **A.** Percent of (S)-FBPMe and (R)-FBPMe: free (white), in site I (green) and in site II (brown). **B** Percent of (R)-FBPMe (●) and (S)-FBPMe (●) in site I of HSA at different $[\text{FBPMe}]/[\text{HSA}]$ ratios.

In summary, the triplet excited state of FBPMe has been used as a chiral reporter for the two binding sites of HSA. The occupation level of the binding sites in HSA has been estimated from regression analysis of the triplet decays at several $[\text{FBPMe}]/[\text{HSA}]$ ratios. A remarkable stereodifferentiation has been found in the triplet lifetimes within the protein microenvironment.

Acknowledgments The UPV (Grant PI 2003-0522 and fellowship to I. V.), the MEC (Grant CTQ2004-03811) and Generalitat Valenciana (Grant GV2004-0536 and Grupos03/082) are gratefully acknowledged for financial support.

References

- (1) He, X. M.; Carter, D. C. *Nature* **1992**, 358, 209.
- (2) Sudlow, G.; Birkett, D. J.; Wade, D. N.; *Mol. Pharmacol.* **1976**, 12, 1052.
- (3) Dockal, M.; Carter, D. C.; Rüker, F. *J. Biol. Chem.* **1999**, 274, 29303.
- (4) Jia, Z.; Ramstad, T.; Zhong, M. *J. Pharm. Biomed. Anal.* **2002**, 405, 413.
- (5) Lucas, L. H.; Price, K. E.; Larive, C. K. *J. Am. Chem. Soc.* **2004**, 126, 14258.
- (6) Evans, A.M. *J. Clin. Pharmacol.* **1996**, 36, 7S.
- (7) Lapique, F.; Muller, N.; Payan, E.; Dubois, N. *Clin. Pharmacokinet.* **1993**, 25, 115.
- (8) Rahman, M. H.; Yamasaki, K.; Shin, Y.-H.; Lin, C. C.; Otagiri, M. *Biol. Pharm. Bull.* **1993**, 16, 1169.
- (9) Bohne, C.; Barra, M.; Boch, R.; Abuin, E. B.; Scaiano, J. C. *J. Photochem. Photobiol. A: Chem.* **1992**, 65, 249.
- (10) Lhiaubet-Vallet, V.; Sarabia, Z.; Bosca, F.; Miranda, M. A. *J. Am. Chem. Soc.* **2004**, 126, 9538.
- (11) Davies, N. M. *Clin. Pharmacokinet.* **1995**, 28, 100.
- (12) Jiménez, M. C.; Miranda, M. A.; Tormos, R.; Vayá, I. *Photochem. Photobiol. Sci.* **2004**, 3, 1038.
- (13) Fehske, K. J.; Müller, K. J.; Schläfer, U.; Wollert, U. *Prog. Drug. Protein Binding*, **1981**, 5-15.



ABSTRACT FOR WEB PUBLICATION

The triplet excited state of flurbiprofen methyl ester (FBPMc) has been used as a chiral reporter for the two binding sites of human serum albumin (HSA). The occupation level of the binding sites has been estimated from regression analysis of the triplet decays at several [FBPMc]/[HSA] ratios. The data agree with two high affinity binding sites (I and II) that are populated to a different extent. A remarkable stereodifferentiation has been found in the drug triplet lifetimes within the protein microenvironment.
